

**Synthesis of polysubstituted pyrroles by the
reaction of enaminoketones, arylglyoxals,
and *N,N*-dimethylbarbituric acid**


Nadezhda N. Kolos^{1*}, Natal'ya V. Chechina¹

¹ *V. N. Karazin Kharkiv National University,
4 Svobody Sq., Kharkiv 61022, Ukraine, e-mail: kolos_n@ukr.net*

SUPPLEMENTARY INFORMATION


^1H , ^{13}C NMR spectra and DEPT-135 spectra of compound

4a




cn64_C13

Automated Probe tuning parameter




cn641_C13dept135
DEPT-135


131.92
130.33
128.46




^1H , ^{13}C NMR spectra and DEPT-135 spectra of compound 4b




cn667_C13
Automated Probe Tuning parameter
ft



cn667_C13dept135
DEPT-135




^1H , ^{13}C NMR spectra and DEPT-135 spectra of compound 4c




cn661_C13


Automated Probe tuning parameter
ft



cn661_C13dept135
DEPT-135




^1H , ^{13}C NMR spectra and DEPT-135 spectra of compound 4d




cn664_C13

Automated Probe tuning parameter


ft



cn664_C13dept135
DEPT-135




^1H , ^{13}C NMR spectra and DEPT-135 spectra of compound 5a



cn697_C13

Automated Peak Picking parameter

ft



cn697_C13dept135
DEPT-135

